

Fall 2007, Vol. 13, No 3

The Prevalence of Latex-Specific IgE in Patients

This Article originally appeared in Fall 2007 issue of the *The Alert* Newsletter, Vol 13, No 3

Volume 40, Issue 4, Pages 411-419 (October 2002)

The prevalence of latex-specific IgE in patients presenting to an urban emergency department original article

Presented at the Society of Academic Emergency Medicine annual meeting, San Francisco, CA, May 2000.

Mary Grzybowski, PhD, MPH, Dennis R. Ownby, MD, Emanuel P. Rivers, MD, MPH, Douglas Ander, MD, Richard M. Nowak, MD

Received 12 November 2001; received in revised form 29 May 2002; accepted 14 June 2002

Abstract

A correction to this article appears on page 160 of the January 2003 issue.

Study objective:

Since the first report of immediate hypersensitivity to latex was documented in 1979, latex allergy has been increasingly recognized as a medical problem in patient populations. However, there are no data available regarding the seroprevalence of latex sensitization in patients presenting to the emergency department. The overall aim of this study was to determine the prevalence of latex IgE seropositivity (L-IgE) in ED patients.

Methods:

We measured latex-specific IgE antibodies among a convenience sample of 1,027 patients using the AlaSTAT assay. We also measured serum IgE antibodies specific for 12 common inhalant allergens using the AlaTOP Microplate Allergy Screen assay. Demographic data were collected. Two questions related to latex allergy were queried. Descriptive statistics are presented. Point estimates and 95% confidence intervals (CIs) were calculated for each seroprevalence test. Odds ratios (ORs) and 95% CIs were used to assess the relationship between sex, race, and seropositivity to inhalant allergens and L-IgE.

Results:

The participation rate was 90% (1,027). The mean age of the patients was 46.9 years, 47.6% (489) were male, and 13.9% (143) were white. Eight-four (8.2%; 95% Ci 6.5% to 10.0%) had positive L-IgE results, and 23.8% (20) of patients with positive L-IgE results were classified as having strongly positive results. Bivariate analyses showed that being nonwhite (OR 4.7, 95% Ci 1.5 to 15.1) and being seropositive for inhalant allergens (OR 7.4; 95% Ci 4.2 to 13.1) were associated with L-IgE.

Conclusion:

The prevalence of latex sensitization in our sampling is substantial and higher than previously estimated in the general adult population. The clinical significance of seropositivity requires further evaluation. [Ann Emerg Med. 2002;40:411-419.]

PII: S0196-0644(02)00063-X

doi:10.1016/S0196-0644(02)00063-X

© 2002 American College of Emergency Physicians. Published by Elsevier Inc. All rights reserved.

This study is the first large-scale study to document the prevalence of latex sensitivity in patients to an ED. The overall percentage of seropositive for latex-specific IgE antibodies was 8.2%.

For reprints or copy of the full article contact:

Mary Grzybowski, PhD, MPH

6G University Health Center
4201 St. Antoine
Detroit, MI 48201

313-745-4347

Email: mgrzyb@med.wayne.edu [1]

Or Visit: <http://www.annemergmed.com/> [2] article/PIIS019606440200063X/ abstract

Reprinted with Permission

Reprinted from

American Latex Allergy Association

P.O. Box 198

Slinger, WI 53086

Phone: 262-677-9707 1-888-97-ALERT

Website: www.latexallergyresources.org

Source URL: <http://latexallergyresources.org/fall-2007-vol-13-no-3>

Links:

[1] <mailto:mgrzyb@med.wayne.edu>

[2] <http://www.annemergmed.com/>